Exhibit 14


. .

Current URL: http://health.clevelandclinic.org/2013/06/the-truth-about-vaginal-mesh/

The Truth About Vaginal Mesh

The story behind sensational headlines

By Women's Health Team | 6/17/13 11:17 a.m.



With so much negative information out there, what are the facts about vaginal mesh?

"Women need to know that they can't believe everything they hear on the news," says gynecologist Marie Paraiso, MD [LINK: http://my.clevelandclinic.org/staff_directory/staff_display.aspx? doctorid=1835], Head, Section of Urogynecology and Reconstructive Pelvic Surgery. "Early meshes were not well designed for acceptance into the body, but not all mesh is bad."

For women who have had vaginal mesh inserted, Dr. Paraiso advises watching for the most common complications. If your doctor is recommending vaginal mesh, be sure he or she is highly skilled for the procedure and that other alternatives have been ruled out.

How vaginal mesh works

Vaginal mesh has been used since about 2005 to treat POP, and is very similar to the mesh products that have been used to treat hernias for many years.

"It's placed in the space between the vagina and bladder or rectum and attached to the ligaments in the pelvis," says Dr. Paraiso. "It spans the area where the prolapse is taking place and gives the vagina the support it once had."

The way these early vaginal meshes were designed, along with the inexperience of many physicians using them, led the Food and Drug Administration to issue a statement in 2008 [LINK: http://www.fda.gov/medicaldevices/safety/alertsandnotices/publichealthnotifications/ucm061976 .htm] warning of serious complications from these devices, she says. A second warning followed in 2011.

What are the complications?

Most complications from vaginal mesh occur in the first three to six months following the procedure, says Dr. Paraiso.

If you've received a vaginal mesh procedure, watch for:

- Pain during intercourse
- Other types of chronic pain
- Fever or vaginal discharge that's bloody or malodorous
- A recurrence of the prolapse

What are the alternatives?

Every woman who receives a recommendation for vaginal mesh should ask about alternative methods of treatment. Dr. Paraiso and her colleagues use vaginal mesh rarely, and only in select cases where it's needed.

Most frequently, she can use a woman's own tissues to create additional support for the pelvic organs.

"There are surgeons with very good experience who are well-prepared for pelvic floor surgery," says Dr. Paraiso. "Ask your doctor what type of training and experience they have in treating POP, and ask what your alternatives are."

More information

Incontinence and Prolapse Treatment Guide [LINK:

http://www.clevelandclinic.org/lp/pelvic_organ_prolapse_incontinence/index.html?
utm campaign=prolapsequide-url&utm medium=offline&utm source=redirect]



We welcome your comments. However, we cannot provide a medical opinion without an in-person consultation. To learn about Cleveland Clinic services available to you, please fill out our WebMail form.



800.801.2273

Home

About Cleveland Clinic

Careers at Cleveland Clinic

Giving

Office of Diversity & Inclusion

Community Outreach

Research & Innovations

Free Health eNewsletters

Resources for Medical Professionals

Media Relations

For Employees

Locations

Main Campus

Cleveland Clinic Children's

Northeast Ohio Family Health Centers

Northeast Ohio Community Hospitals

Northeast Ohio Specialty Centers

Florida

Canada

Nevada

Abu Dhabi

Site Information

Send Us Feedback

Copyright, Reprints & Licensing

Site Map

About this Website

Website Terms of Use

Website Privacy Policy

Notice of Privacy Practices

Need Help Paying Your Medical Bills?



We comply with the HONcode standard for trustworthy health information: verify here.



Cleveland Clinic © 1995-2014. All Rights Reserved. 9500 Euclid Avenue, Cleveland, Ohio 44195 | 800.223.2273 | TTY 216.444.0261